

# COLEMAN'S $\mathcal{L}$ -INVARIANT AND FAMILIES OF MODULAR FORMS

GLENN STEVENS

## CONTENTS

Statement of results	1
1. The Gauss-Manin connection with Frobenius structure	3
2. Coleman's $\mathcal{L}$ -invariant.	4
3. Families of modular forms	6
4. Some Pairings	8
5. Proof of Theorem B	9
References	12

## STATEMENT OF RESULTS

Let  $p$  be a prime  $> 2$  and  $N$  be a positive integer with  $p \nmid N$ . Let  $f$  be a classical newform over  $\Gamma_0(Np)$  of even weight  $k_0 + 2 \geq 2$  and assume  $f$  is split multiplicative at  $p$ , thus

$$a_p(f) = p^{k_0/2}$$

where  $a_p(f)$  is the eigenvalue of the  $U$ -operator at  $p$  acting on  $f$ . Under these hypotheses, Coleman [3] defined an  $\mathcal{L}$ -invariant  $\mathcal{L}(f)$  which he conjectured to be equal to the higher weight Mazur-Tate-Teitelbaum  $\mathcal{L}$ -invariant [16]. In this paper we will prove Coleman's conjecture. More precisely, let  $\mathcal{X} := \mathbb{Z}/(p-1)\mathbb{Z} \times \mathbb{Z}_p$  with  $\mathbb{Z}$  embedded in  $\mathcal{X}$  diagonally and let  $L_p(f, -) : \mathcal{X} \rightarrow \mathbb{C}_p$  be the  $p$ -adic  $L$ -function attached to  $f$  as in [16]. We will prove the following theorem.

**Main Theorem.**  $L'_p(f, 1 + k_0/2) = \mathcal{L}(f) \cdot L_\infty(f, 1 + k_0/2)$ .

In the special case of weight two ( $k_0 = 0$ ), in which case,  $\mathcal{L}(f)$  takes the familiar form  $\mathcal{L}(f) = \log(q_f)/\text{ord}(q_f)$  when  $f$  has rational fourier coefficients, this was conjectured by Mazur, Tate, and Teitelbaum [16] and proved by Ralph Greenberg and the author in [11, 12]. In case  $f$  is split multiplicative of weight  $k_0 + 2 > 2$ , Mazur, Tate, and Teitelbaum offered no precise formula for

---

Research partially supported by NSF grants DMS 9401553, DMS 9701782, and DMS 0071065.

$\mathcal{L}(f)$ , but they did predict that  $\mathcal{L}(f)$  could be described purely in terms of local  $p$ -adic data associated to  $f$  and that, in particular,  $\mathcal{L}(f)$  should not change when  $f$  is twisted by a Dirichlet character  $\chi$  with  $\chi(p) = 1$ . Three separate and apparently independent definitions of  $\mathcal{L}(f)$  were later proposed. The first was given by Jeremy Teitelbaum [18], but only in the case where  $f$  corresponds to a quaternionic modular form via the Jacquet-Langlands correspondence. Robert Coleman gave an analogous definition in [3] in the general case, which we will briefly recall in section 2 of this paper. A third definition was proposed by Fontaine and Mazur [15] based on Fontaine's theory of semistable  $p$ -adic Galois representations. These three definitions gave rise to three separate conjectures of Mazur-Tate-Teitelbaum type. All three of these conjectures have now been proved.

The  $\mathcal{L}$ -invariants of Coleman and Teitelbaum can be approximated  $p$ -adically on a computer, which enabled early numerical confirmation of the Coleman and Teitelbaum conjectures [6, 7, 18]. On the other hand, the Fontaine-Mazur  $\mathcal{L}$ -invariant appears to be beyond the reach of a computer. Nevertheless, it was the Fontaine-Mazur version of the conjecture that was the first to be proved – in 1996, by Kato, Kurihara, and Tsuji [13]. The Coleman version of the conjecture was established by the author shortly thereafter and described in [17], thus also proving indirectly that the Fontaine-Mazur and Coleman  $\mathcal{L}$ -invariants are the same. Coleman and Iovita [5] later gave a direct proof that *all three*  $\mathcal{L}$ -invariants—including Teitelbaum's invariant when it is defined—are the same. For an excellent overview of the history of the  $\mathcal{L}$ -invariant and the Mazur-Tate-Teitelbaum conjecture, see Colmez's survey [10]. The connection with Kato's Euler systems and the  $p$ -adic Birch-Swinnerton-Dyer conjecture, including the proof by Kato, Kurihara, and Tsuji given in the language of  $(\varphi, \Gamma)$ -modules, is also beautifully described in Colmez's Bourbaki seminar notes [9].

As in the weight two case (see [11, 12]), our proof of Coleman's conjecture in the higher weight case divides naturally into two steps (Theorems A and B below). To state Theorems A and B, we first recall that Coleman [2] constructed a  $p$ -adic analytic family  $f_k$  of overconvergent  $p$ -adic modular forms passing through our fixed newform  $f$ . This family is defined for  $k$  in an open set  $B \subseteq \mathcal{X}$  containing  $k_0$  and satisfies  $f_{k_0} = f$ . Coleman's family is an eigenfamily for the  $U$ -operator and we may therefore consider the eigenvalue  $\alpha(k)$  of  $U$  acting on  $f_k$ . The function  $\alpha(k)$  is a  $p$ -adic analytic function of  $k \in B$  so we may consider the derivative of  $\alpha$  at the special point  $k_0 \in B$ .

**Theorem A.**  $L'_p(f, 1 + k_0/2) = -2 \cdot p^{-k_0/2} \cdot \alpha'(k_0) \cdot L_\infty(f, 1 + k_0/2)$ .

Just as in the weight two case, the proof of Theorem A depends on the existence of a two variable  $p$ -adic  $L$ -function with certain properties. The existence of such a  $p$ -adic  $L$ -function was proved

in the higher weight case in [17]. With the two-variable  $p$ -adic  $L$ -function in hand, the proof of theorem A proceeds exactly as in the weight two case (see [11, 12]).

The rest of this note is dedicated to proving the following theorem.

**Theorem B.**  $\mathcal{L}(f) = -2 \cdot p^{-k_0/2} \cdot \alpha'(k_0)$ .

The Main Theorem is an immediate consequence of Theorems A and B. We remark that Colmez [8] has also proven Theorem B, but in terms of the Fontaine-Mazur  $\mathcal{L}$ -invariant.

## 1. THE GAUSS-MANIN CONNECTION WITH FROBENIUS STRUCTURE

We adopt Coleman's notations as in [3] with only one modification. Namely, we will add full level 2 structure to the moduli space. This rigidifies the setup and simplifies the calculations (see especially the proof of Proposition 3.1(2)). We let  $X$  be the modular curve  $X(Np, 2)$  with level  $Np$  structure (a cyclic subgroup of order  $Np$ ) plus full level 2 structure. (If  $2|N$  we assume that the additional level 2 structure extends the 2-part of the level  $N$  structure.) The  $p$ -adic rigid analytic space  $X^{an}$  attached to  $X$  is the union of three disjoint parts, namely,

$$X^{an} = Z_\infty \cup W \cup Z_0$$

where  $Z_\infty$  and  $Z_0$  are the ordinary affinoids containing the  $\infty$  and 0-cusps respectively, and  $W$  is the union of the supersingular annuli. Following Coleman, we write  $W_\infty = Z_\infty \cup W$  and  $W_0 = Z_0 \cup W$ .

Let  $Y = Y(Np, 2)$  denote  $X$  with the cusps deleted. Let  $\pi : E \rightarrow Y$  be the universal elliptic curve with level structure over  $Y$  and let  $\mathcal{H}$  be the relative de Rham cohomology sheaf over  $X$  with log singularities at the cusps. Then  $\mathcal{H}$  is a coherent  $\mathcal{O}$ -module locally free of rank 2 over  $X$ . As Katz explains in [14] we have a canonical decomposition

$$\mathcal{H} = \underline{\omega}^{-1} \oplus \underline{\omega}$$

where  $\underline{\omega} := \pi_* \Omega_{E/Y}^1$ . For any nonnegative integer  $k$  we let

$$\mathcal{H}_k := \text{Sym}^k(\mathcal{H}) = \underline{\omega}^{-k} \oplus \underline{\omega}^{2-k} \oplus \dots \oplus \underline{\omega}^k.$$

The Gauss-Manin connection  $\nabla : \mathcal{H} \rightarrow \mathcal{H} \otimes \Omega$  induces a connection

$$\nabla : \mathcal{H}_k \rightarrow \mathcal{H}_k \otimes \Omega$$

for each integer  $k \geq 0$ , which we also call the Gauss-Manin connection.

The Deligne-Tate map ([14]) preserves  $Z_\infty$  and extends to a wide open neighborhood of  $Z_\infty$  *properly* contained in  $W_\infty$ . Accordingly, the Gauss-Manin connection is endowed with a natural Frobenius structure over some sufficiently small wide open neighborhood of  $Z_\infty$ . Katz spells out

precisely how big this neighborhood can be, but this is a technical point that we will not need. It will be convenient to simplify the notation and write  $Z_\infty^\dagger$  to denote a choice of such a wide open neighborhood of  $Z_\infty$  with the additional property that the intersection of  $Z_\infty^\dagger$  with any supersingular annulus is a concentric subannulus.

For  $k$  an integer, let  $M_{k+2}^\dagger := \underline{\omega}^{k+2}(Z_\infty^\dagger)$  denote the space of overconvergent  $p$ -adic modular forms of weight  $k+2$  and level  $(Np, 2)$  as before. For  $k \geq 0$  we let

$$\kappa : M_{k+2}^\dagger \longrightarrow \mathcal{H}_k \otimes \Omega(Z_\infty^\dagger)$$

be the Kodaira Spencer map (see §4 of [1]). The canonical projection  $\mathcal{H}_k \longrightarrow \underline{\omega}^{-k}$  induces a surjection  $\mathcal{H}_k(Z_\infty^\dagger) \longrightarrow M_{-k}^\dagger$ , and Coleman proves in [1] that there is a canonical  $\mathbb{Q}_p$ -linear section

$$\nu : M_{-k}^\dagger \longrightarrow \mathcal{H}_k(Z_\infty^\dagger)$$

satisfying the equation

$$\nabla(\nu(g)) = \kappa(\theta^{k+1}g)/k! \in \mathcal{H}_k \otimes \Omega(Z_\infty^\dagger)$$

for any  $g \in M_{-k}^\dagger$ . Here  $\theta^{k+1} : M_{-k}^\dagger \longrightarrow M_{k+2}^\dagger$  is the operator defined on  $q$ -expansions by

$$\theta^{k+1} : \sum_{n \geq 0} a_n q^n \longmapsto \sum_{n \geq 0} n^{k+1} a_n q^n.$$

For details, see Proposition 4.3 of [1].

Following Katz [14], Coleman [3] also defines a Frobenius structure on  $Z_\infty^\dagger$  which gives rise to a ‘‘Frobenius operator’’  $\Phi$  acting on the cohomology of  $\mathcal{H}_k$ ,  $\underline{\omega}^k$ , and  $\Omega$ . Moreover,  $\Phi$  commutes with  $\nabla : \mathcal{H}_k \longrightarrow \mathcal{H}_k \otimes \Omega$  on  $Z_\infty^\dagger$  (see §11 of [3]). On  $q$ -expansions of modular forms of weight  $k$ ,  $\Phi$  is given by  $\Phi = p^k V$  where  $V$  is the operator on  $q$ -expansions given by  $V(f)(q) = f(q^p)$ , i.e.

$$V : \sum_{n \geq 0} a_n q^n \longmapsto \sum_{n \geq 0} a_n q^{np},$$

## 2. COLEMAN’S $\mathcal{L}$ -INVARIANT.

In this section we recall Coleman’s definition of the  $\mathcal{L}$ -invariant  $\mathcal{L}(f)$  of a split multiplicative  $p$ -newform  $f$  of weight  $k_0 + 2 \geq 2$ . Let  $\mathcal{H}_{k_0}^*$  denote the complex of sheaves associated to  $\mathcal{H}_{k_0} \xrightarrow{\nabla} \mathcal{H}_{k_0} \otimes \Omega$  and consider the hypercohomology  $\mathbb{H}^1(X, \mathcal{H}_{k_0}^*)$  with respect to the covering  $\{W_\infty, W_0\}$  of  $X$ . The Hecke operators act on this space and the systems of eigenvalues that occur in it are the same as those that occur in the space of classical modular forms of weight  $k_0$  and corresponding level. In particular, letting  $K$  be the field generated over  $\mathbb{Q}_p$  by the eigenvalues of the Hecke operators acting on  $f$ , we obtain a  $\mathbb{Q}_p$ -subspace  $H(f) \subseteq \mathbb{H}^1(X, \mathcal{H}_{k_0}^*)$  endowed with an action of the field  $K$  with the property that  $H(f)$  is a 2-dimensional  $K$ -vector space on which the Hecke

operators act as scalars according to the eigenvalues of  $f$ . Using his theory of  $p$ -adic integration, Coleman endows  $H(f)$  with a natural monodromy module structure in which the monodromy is *non-trivial*. In [15], Mazur attaches an  $\mathcal{L}$ -invariant to any two dimensional monodromy module with non-trivial monodromy. Coleman's  $\mathcal{L}$ -invariant can be defined simply as the  $\mathcal{L}$ -invariant of Coleman's monodromy module.

We will use the more concrete definition that Coleman gives in [3]. For simplicity, we assume  $k_0 > 0$  so that there are no nonzero horizontal sections of  $\mathcal{H}_{k_0}$  defined on all of  $W_\infty$  nor on all of  $W_0$ , i.e.  $H^0(W_\infty, \mathcal{H}_{k_0}^*) = H^0(W_0, \mathcal{H}_{k_0}^*) = 0$ . On the other hand, one generally does find non-zero horizontal sections of  $\mathcal{H}_{k_0}$  on the supersingular annuli  $W = W_\infty \cap W_0$ . Indeed, Coleman constructs two maps

$$\sigma, \rho : M_{k_0+2} \longrightarrow H^0(W, \mathcal{H}_{k_0}^*)$$

defined on the space  $M_{k_0+2}$  of classical modular forms of weight  $k_0 + 2$  and appropriate level. The map  $\sigma$  is defined using Coleman integration (Definition 2.1 below) while the map  $\rho$  is defined in terms of residues (Definition 2.2).

Let  $k \geq 0$  and  $f \in M_{k+2}$  be a classical Hecke eigenform. Let  $\alpha$  be the eigenvalue of the  $U$ -operator acting on  $f$ . We suppose  $\alpha \neq 0$ . The differential form  $\omega_f := \kappa(f) \in \mathcal{H}_k \otimes \Omega(W_\infty)$  represents a cohomology class  $[\omega_f] \in H^1(W_\infty, \mathcal{H}_k)$  and the Frobenius operator  $\Phi$  acts on  $\omega_f$  and also on  $[\omega_f]$ . Indeed, we have  $\Phi([\omega_f]) = \frac{p^{k+1}}{\alpha} \cdot [\omega_f]$ . Now Coleman's integration theory gives us a well-defined flabby antiderivative  $I_\infty(f)$  defined on all of  $W_\infty$  which is rigid analytic on the ordinary residue disks, is log-analytic on the supersingular annuli and satisfies the following two properties

- $I_\infty(f)$  satisfies the differential equation

$$\nabla(I_\infty(f)) = \omega_f \quad \text{on } W_\infty.$$

- the flabby analytic section

$$I_\infty(f) - \frac{\alpha}{p^{k+1}} \Phi(I_\infty(f))$$

of  $\mathcal{H}_k$  is rigid analytic on  $Z_\infty^\dagger$  (i.e. not only on  $Z_\infty$ , but also on some wide open neighborhood of  $Z_\infty$ ).

Similar considerations give rise to a well-defined flabby analytic section  $I_0(f)$  of  $\mathcal{H}_k$  over  $W_0$  satisfying the differential equation

$$\nabla(I_0(f)) = \omega_f \quad \text{on } W_0.$$

Both  $I_0(f)$  and  $I_\infty(f)$  are defined on the overlap  $W = W_\infty \cap W_0$ . Coleman makes the following definition.

**Definition 2.1.** *If  $f \in M_{k+2}$  is a classical Hecke eigenform then we define  $\sigma(f) \in H^0(W, \mathcal{H}_k^*)$  to be the horizontal section of  $\mathcal{H}_k$  on  $W$  given by*

$$\sigma(f) := I_\infty(f)|_W - I_0(f)|_W.$$

The residue map  $\rho : M_{k+2} \longrightarrow H^0(W, \mathcal{H}_k^*)$  is defined using the map

$$\text{Res} : \mathcal{H}_k \otimes \Omega(Z_\infty^\dagger) \longrightarrow H^0(W, \mathcal{H}_k^*)$$

which in turn is defined by  $\text{Res}(\omega) :=$  the unique horizontal section of  $\mathcal{H}_k$  on  $W$  whose restriction to  $Z_\infty^\dagger \cap W$  is the residue of  $\omega$  restricted to this disjoint union of oriented annuli. Note that here as elsewhere we use the standard orientation of the annuli, i.e. the orientation in which  $Z_\infty$  is *interior* to  $W$ .

**Definition 2.2.** *Given  $f \in M_{k+2}^\dagger$ , we let  $\omega_f := \kappa(f) \in \mathcal{H}_k \times \Omega(Z_\infty^\dagger)$  and define  $\rho(f) := \text{Res}(\omega_f)$ .*

**Definition 2.3.** *Coleman's  $\mathcal{L}$ -invariant of a split multiplicative newform  $f \in M_{k+2}$  is defined to be the unique element  $\mathcal{L}(f) \in K$  for which  $\sigma(f) = \mathcal{L}(f) \cdot \rho(f)$ .*

The existence and uniqueness of such an  $\mathcal{L}$ -invariant was, of course, proved by Coleman (see [3]).

### 3. FAMILIES OF MODULAR FORMS

First of all we have the Eisenstein family. For each integer  $k$  there is an overconvergent  $p$ -adic modular form  $E_k$  of weight  $k$  whose  $q$ -expansion is given by

$$E_k := 1 + 2\zeta_p(1-k)^{-1} \sum_{k \geq 1} \sigma_{k+1}^*(n) q^n.$$

Here  $\zeta_p(s)$  is the Kubota–Leopoldt  $p$ -adic zeta function and when  $k = 0$  the above equality is understood to mean  $E_0 = 1$ . (Recall  $\zeta_p(s)$  has a simple pole at  $s = 1$ ). For integral  $k \geq 0$  we set

$$t_k := \frac{1}{2}\zeta_p(1+k) \cdot E_{-k} \quad \text{and} \quad G_k := \frac{1}{2}\zeta_p(-1-k) \cdot E_{k+2}.$$

Then  $t_k \in M_{-k}^\dagger$  is an overconvergent modular form of weight  $-k$  and  $G_k \in M_{k+2}$  is a classical modular form of weight  $k+2$ . The family  $t_k$  extends to a meromorphic family of Eisenstein series for  $k \in \mathcal{X}$  with a simple pole at  $k = 0$  and  $G_k$  defines a meromorphic family with a simple pole at  $k = -2$ . Moreover  $G_k = t_{-2-k}$ . The special point  $k = 0$  will play a crucial role in the proof of Theorem B.

**Proposition 3.1.** *The following assertions hold.*

(1) *The family  $t_k$ ,  $k \in \mathcal{X}$ , has a simple pole at  $k = 0$  with residue given by*

$$\lim_{k \rightarrow 0} kt_k = \frac{1}{2} \cdot \left(1 - \frac{1}{p}\right).$$

(2) *The residue of  $G_0$  along any supersingular annulus is  $1/2$ :*

$$\rho(G_0) = -\frac{1}{2}.$$

**Proof.** The first assertion is an immediate consequence of the well-known fact that the Kubota-Leopoldt  $p$ -adic zeta function  $\zeta_p(s)$  has a simple pole at  $s = 1$  and that the residue at  $s = 1$  is given by

$$\lim_{s \rightarrow 1} (s - 1)\zeta_p(s) = \left(1 - \frac{1}{p}\right).$$

To prove the second assertion, we first consider the special case  $N = 1$ . Then  $\eta = \kappa(G_0)$  is a section of  $\Omega$  over  $Y$  which extends to a meromorphic section over  $X$  with simple poles along the cusps. We want to compute  $\text{Res}(\eta) \in H^0(W)$ . We remark first of all that since the eigenvalues of the Hecke operators acting on  $\eta$  are known, they are also known on  $\text{Res}(\eta)$ . Indeed, the eigenvalues are the same as those acting on constant functions on  $W$ . Hence  $\text{Res}(\eta)$  is a constant. To determine what the constant is we use the fact that the sum of the residues along the oriented annuli at the “edges” of  $W_\infty \setminus \{\text{cusps}\}$  is equal to zero. For the supersingular annuli this is the orientation we used above to define  $\rho(f)$ . However, the annuli around the cusps are orientated so that the cusps are *exterior* to the annuli, which is opposite the one used to compute the constant terms of Eisenstein series.

Under our assumption  $N = 1$ , there are a total of three cusps in  $W_\infty$  corresponding to the three cusps of  $X(2)$ . The constant terms of  $G_0$  are the same at all of these cusps since  $G_0$  is modular of level  $p$ . Since the natural map  $X \rightarrow X_0(p)$  is ramified of order 2 at each of these cusps and since the constant term of  $G_0$  at the infinity cusp is  $(1 - p)/24$  we conclude that the sum of the residues along the oriented annuli around the cusps is  $(p - 1)/4$ . Hence the sum of the residues along the supersingular annuli is  $(1 - p)/4$ . But a simple calculation shows that the number of supersingular annuli in  $X$  is  $(p - 1)/2$ . Hence the residue along any supersingular annulus is  $-1/2$ . This proves (2) when  $N = 1$ .

The general case follows at once since for arbitrary  $N$ , the map  $X(Np, 2) \rightarrow X(p, 2)$  is unramified over the supersingular annuli. This completes the proof of the proposition.  $\square$

We can remove Euler factors at  $p$  using the operator  $V$  on overconvergent modular forms defined on  $q$ -expansions by the formula  $V(f)(q) = f(q^p)$ . If  $F$  is an eigenform, then we let  $F^0$  denote the

eigenform obtained by removing the Euler factor at  $p$ . Thus, we have the families

$$\begin{aligned} t_k^0 &:= t_k - V(t_k) \\ G_k^0 &:= G_k - V(G_k) \\ f_k^0 &:= f_k - \alpha(k)V(f_k) \end{aligned}$$

For  $k \geq 0$  we let  $\eta_k := \kappa(G_k)$  and  $\eta_k^0 := \kappa(G_k^0)$  where  $\kappa : \mathcal{M}_{k+2} \longrightarrow \mathcal{H}_k \otimes \Omega$  is the Kodaira-Spencer map. We also set  $g_k := \nu(t_k)$  and  $g_k^0 := \nu(t_k^0)$ . Then since  $\theta^{k+1}t_k^0 = G_k^0$  it follows that

$$\nabla(g_k^0) = G_k^0.$$

Finally, for each integer  $k \geq 0$  we may let  $s_k := I_\infty(f_k)$  be the Coleman integral of  $f_k$  defined in section 1. Then  $s_k$  is a flabby section of  $\mathcal{H}_k$  over  $W_\infty$ . This section is characterized by the property that

$$s_k^0 := s_k - \frac{\alpha(k)}{p^{k+1}} \cdot \Phi(s_k)$$

is a rigid analytic section of  $\mathcal{H}_k$  over  $Z_\infty^\dagger$ . Hence there is an overconvergent modular form  $\phi_k^0 \in M_{-k}^\dagger$  such that

$$\nu(\phi_k^0) = s_k^0.$$

Hence  $\theta^{k+1}(\phi_k^0) = f_k^0$ . Finally, set

$$\begin{aligned} \omega_k &:= \kappa(f_k), \\ \omega_k^0 &:= \kappa(f_k^0). \end{aligned}$$

#### 4. SOME PAIRINGS

As in the introduction, we fix an integer  $k_0 \geq 0$ . For each integer  $k \geq 0$  cup product on the de Rham cohomology of the fibers of  $E/X$  induces a natural pairing  $[\cdot, \cdot] : \mathcal{H}_k \times \mathcal{H}_{k+k_0} \longrightarrow \mathcal{H}_{k_0}$ . This pairing induces natural pairings

$$\begin{aligned} [\cdot, \cdot] &: \mathcal{H}_k \times \mathcal{H}_{k+k_0} \otimes \Omega \longrightarrow \mathcal{H}_{k_0} \otimes \Omega; \\ [\cdot, \cdot] &: \mathcal{H}_k \otimes \Omega \times \mathcal{H}_{k+k_0} \longrightarrow \mathcal{H}_{k_0} \otimes \Omega. \end{aligned}$$

**Proposition 4.1.** *These pairings satisfy the following identity for all  $x \in \mathcal{H}_k$ , and  $y \in \mathcal{H}_{k+k_0}$*

$$\nabla[x, y] = [x, \nabla y] + [\nabla x, y].$$

**Proof.** This follows from the product formula for differentiation. □

We will use a superscript  $\dagger$  to denote overconvergent sections of a sheaf. For example,  $\mathcal{H}_k^\dagger := \mathcal{H}_k(Z_\infty^\dagger)$ . We may then define pairings

$$\begin{aligned} \langle \cdot, \cdot \rangle &: \mathcal{H}_k^\dagger \times \mathcal{H}_{k+k_0}^\dagger \otimes \Omega^\dagger \longrightarrow H^0(W, \mathcal{H}_{k_0}^*) \\ \langle \cdot, \cdot \rangle &: \mathcal{H}_k^\dagger \otimes \Omega^\dagger \times \mathcal{H}_{k+k_0}^\dagger \longrightarrow H^0(W, \mathcal{H}_{k_0}^*) \end{aligned}$$



by defining  $\langle x, y \rangle := \text{Res}([x, y])$  where  $\text{Res} : \mathcal{H}_{k_0}^\dagger \longrightarrow H^0(W, \mathcal{H}_{k_0}^*)$  is the residue map.

Recall the Hecke operators  $U := U_p$  and  $w$  from §8 of [3]. On  $q$ -expansions of modular forms  $U$  is given by

$$U : \sum_{n \geq 0} a_n q^n \longmapsto \sum_{n \geq 0} a_{np} q^n.$$

The operator  $w$  acts on  $\mathcal{H}_{k_0}(W)$  and satisfies  $w^2 = p^{k_0}$  on this space. Hence

$$W_p := p^{-k_0/2} w$$

acts as an involution. Moreover, from §11 of [3] we have  $\Phi = w$  on horizontal sections of  $\mathcal{H}_{k_0}$  over  $W$ , hence  $\Phi = p^{k_0/2} W_p$  on  $H^0(W, \mathcal{H}_{k_0}^*)$ .

**Proposition 4.2.** *Let  $k, k_0$  be non-negative integers and let  $g \in M_{-k}^\dagger$ ,  $f \in M_{k+k_0}^\dagger$ , and  $h \in M_{k+2}^\dagger$ . Then the following assertions hold.*

- (1) For  $x = \nu(g) \in \mathcal{H}_k^\dagger$  and  $\omega = \kappa(f) \in \mathcal{H}_{k+k_0}^\dagger \otimes \Omega^\dagger$  we have  $\langle x, \Phi(\omega) \rangle = p^{k + \frac{k_0}{2} + 1} \cdot W_p(\langle U(x), \omega \rangle)$ ;
- (2) For  $\eta = \kappa(h) \in \mathcal{H}_k^\dagger \otimes \Omega^\dagger$  and  $y \in \mathcal{H}_{k+k_0}^\dagger$  we have  $\langle \eta, \Phi(y) \rangle = p^{k + \frac{k_0}{2}} \cdot W_p(\langle U(\eta), y \rangle)$ .

*Proof.* Since  $p^{k_0/2} W_p = \Phi$  and  $p \cdot \Phi \circ \text{Res} = \text{Res} \circ \Phi$  we have

$$\begin{aligned} p^{k + \frac{k_0}{2} + 1} \cdot W_p(\langle U(x), \omega \rangle) &= p^k \cdot \langle \Phi U(x), \Phi \omega \rangle \\ &= p^k \rho(\Phi U(x) \cdot \Phi \omega) \\ &= \rho(g \cdot \Phi(f)) \\ &= \langle x, \Phi(\omega) \rangle. \end{aligned}$$

This proves (1) and (2) is proved similarly. □

## 5. PROOF OF THEOREM B

The operator  $W_p$  is an involution on  $H^0(W, \mathcal{H}_{k_0})$ . We let superscript  $+$  denote projection to the  $+$ -component under the action of  $W_p$ . Consider the function  $\psi : \mathcal{X} \longrightarrow H^0(W, \mathcal{H}_{k_0})^+$  defined by

$$\psi(k) := \rho(t_k^0 f_{k+k_0}^0)^+ \in H^0(W, \mathcal{H}_{k_0})^+.$$

Since  $t_k^0 f_{k+k_0}^0$  is an analytic family of overconvergent modular forms of weight  $k_0$  we see at once that  $\psi(k)$  is an analytic function of  $k$  defined on a neighborhood of 0 in  $\mathcal{X}$ . For the proof of Theorem B we will calculate  $\psi(0)$  in two ways. First, by direct calculation we express  $\psi(0)$  in terms of  $\rho(f)$ . Then we apply the product rule (Proposition 2) to express  $\psi(0)$  in terms of  $\sigma(f)$ . Comparing these two expressions, Theorem B follows.

Define  $u(k) := p^{-k_0/2} \cdot \alpha(k)$ , the “unit part” of  $\alpha(k)$ .

**Lemma 5.1.** *We have*

$$\psi(0) = -\frac{1}{2} \cdot \left(1 - \frac{1}{p}\right) \cdot u'(k_0) \cdot \rho(f).$$

**Proof.** For an arbitrary integer  $k \geq 0$  we have

$$\psi(k) = \rho(t_k^0 f_{k+k_0}^0)^+ = \langle g_k^0, \omega_{k+k_0}^0 \rangle^+.$$

We also have

$$\begin{aligned} \langle g_k^0, \omega_{k+k_0}^0 \rangle &= \langle g_k, \omega_{k+k_0}^0 \rangle \\ &= \left\langle g_k, \omega_{k+k_0} - \frac{\alpha(k+k_0)}{p^{k+k_0+1}} \Phi(\omega_{k+k_0}) \right\rangle \\ &= \langle g_k, \omega_{k+k_0} \rangle - \frac{\alpha(k+k_0)}{p^{k_0/2}} W_p(\langle U(g_k), \omega_{k+k_0} \rangle) \\ &= \langle g_k, \omega_{k+k_0} \rangle - u(k+k_0) \cdot W_p(\langle g_k, \omega_{k+k_0} \rangle). \end{aligned}$$

The first equality above follows from three facts: (1)  $g_k^0 - g_k$  is in the image of  $\Phi$ ; (2)  $\omega_{k+k_0}^0$  is in the kernel of  $U$ ; and (3) the image of  $\Phi$  is perpendicular to the kernel of  $U$  by Proposition 4.2. The third equality follows from Proposition 4.2(1). The last equality above follows from the fact that the Eisenstein series  $t_k$  is an eigenform for the  $U$ -operator with eigenvalue 1, hence  $U(g_k) = g_k$ .

Now project the above identity to the  $+$ -component for  $W_p$  to get

$$\begin{aligned} \psi(k) &= (1 - u(k+k_0)) \cdot \langle g_k, \omega_{k+k_0} \rangle^+ \\ &= \frac{1 - u(k+k_0)}{k} \cdot \rho(kt_k f_{k+k_0})^+. \end{aligned}$$

Letting  $k \rightarrow 0$ , using Propostion 3.1(1), and noting that  $\rho(f)^+ = \rho(f)$  we obtain

$$\psi(0) = -\frac{1}{2} \cdot \left(1 - \frac{1}{p}\right) \cdot u'(k_0) \cdot \rho(f)$$

and the lemma is proved.  $\square$

Let  $C_\infty := Z_\infty^\dagger \setminus Z_\infty$ . Then  $C_\infty$  is a union of concentric annuli in the supersingular annuli. Note that the pairings  $\langle x, y \rangle$  are well-defined so long as  $x, y$  are rigid on  $C_\infty$ . In particular we have a well-defined pairing

$$\langle \cdot, \cdot \rangle : \Omega^1(C_\infty) \times \mathcal{H}_{k_0}(C_\infty) \longrightarrow \mathcal{H}_{k_0}(W)^\nabla.$$

defined by  $\langle \omega, h \rangle = \text{Res}_W(h\omega)$ , where this latter is defined to be the unique horizontal section on  $W$  extending  $\text{Res}_{C_\infty}(h\omega)$ .

We now turn to an application of Coleman's integration theory. In what follows, we will write a subscript *flog* to denote flabby log-analytic sections of a rigid analytic sheaf. Such sections are, by definition, rigid analytic on the residue disks in the ordinary part of  $X$  and are log-analytic on the supersingular annuli. For details, see §10 of [3] and also [4].

**Lemma 5.2.** *Let  $e \in \mathcal{O}_{f\log}(W_\infty)$  be any Coleman integral of  $\eta_0$  (well-defined up to a constant). Restrict  $e$  to the supersingular annuli  $W$  and let  $h = e - W_p(e) \in \mathcal{O}_{\log}(W)$ . Let  $z = h \cdot \rho(f) \in \mathcal{H}_{k_0, \log}(W)$ , and let  $z^0 := z - p^{-1-k_0/2}\Phi(z) \in \mathcal{H}_{k_0, \log}(C_\infty)$ . Then  $z, z^0$  have the following properties.*

- (1)  $z^0$  is rigid on  $C_\infty$ .
- (2)  $s_{k_0} + z$  is rigid on  $W$ .
- (3)  $\langle \eta_0, z^0 \rangle = 0$ .
- (4)  $W_p(z) + z = 0$  on the supersingular annuli  $W$ .

**Proof.** (1) Since  $e$  is a Coleman integral of  $\eta_0$ , we have  $e^0 := e - p^{-1}\Phi(e)$  is rigid on  $Z_\infty^\dagger$ . Since  $W_p(\eta_0) = -\eta_0$  on  $X$ , we have  $W_p(e) + e$  is constant on  $W$ . It follows that  $h^0 := h - p^{-1}\Phi(h)$  is also rigid on  $C_\infty$ . On the other hand,  $\Phi(\rho(f)) = p^{k_0/2}\rho(f)$ . Hence  $z^0 = h^0 \cdot \rho(f)$ , which is rigid on  $C_\infty$ .

(2) By definition,  $\nabla(s_{k_0}) = \kappa(f)$ . Hence,  $\text{Res}_W(\nabla(s_{k_0})) = \rho(f)$ . On the other hand,  $\text{Res}_W(\nabla(z)) = \text{Res}_W(dh) \cdot \rho(f)$ . But  $dh = 2\eta_0$  and we have shown in Proposition 3.1 that  $\text{Res}_W(\eta_0) = -1/2$ , hence  $\text{Res}_W(\nabla(z)) = \rho(f)$ . We therefore have  $\text{Res}_W(\nabla(s_{k_0} + z)) = 0$  and it follows that  $s_{k_0} + z$  is rigid on  $W$ , as claimed.

(3) We have  $\langle \eta_0, z^0 \rangle = \langle \eta_0, h^0 \rangle \cdot \rho(f)$ . Moreover,  $\langle \eta_0, h^0 \rangle = \langle \eta_0^0, h^0 \rangle$  because the image of  $\Phi$  is orthogonal to the kernel of  $U$ . But,  $\langle \eta_0^0, h^0 \rangle = \text{Res}_W(h^0 \eta_0^0) = \frac{1}{2} \text{Res}_W(h^0 dh^0) = 0$ , since  $h^0 dh^0$  is an exact differential on  $C_\infty$ .

(4) Since  $W_p(\rho(f)) = \rho(f)$ , this follows immediately from the definition of  $z$ .

This completes the proof of Lemma 5.2. □

**Lemma 5.3.**  $\psi(0) = \frac{1}{4} \left(1 - \frac{1}{p}\right) \sigma(f)$ .

**Proof.** As in the first line of the proof of Lemma 5.1 we have

$$\psi(0) = \langle g_0^0, \omega_{k_0}^0 \rangle^+.$$

But  $\omega_{k_0}^0$  is an exact differential, indeed  $\nabla s_{k_0}^0 = \omega_{k_0}^0$ . Moreover,  $\nabla g_0^0 = \eta_0^0$ . Hence, by Proposition 4.1 we have

$$\nabla[g_0^0, s_{k_0}^0] = [\eta_0^0, s_{k_0}^0] + [g_0^0, \omega_{k_0}^0].$$

Taking residues of both sides of this equality along the supersingular annuli we obtain

$$0 = \langle \eta_0^0, s_{k_0}^0 \rangle + \langle g_0^0, \omega_{k_0}^0 \rangle.$$

Hence  $\psi(0) = -\langle \eta_0^0, s_{k_0}^0 \rangle^+$ , which we calculate as follows:

$$\begin{aligned}
\langle \eta_0^0, s_{k_0}^0 \rangle &= \langle \eta_0, s_{k_0}^0 \rangle \\
&= \langle \eta_0, s_{k_0}^0 + z^0 \rangle \\
&= \langle \eta_0, (s_{k_0} + z) - \frac{1}{p^{k_0/2+1}} \cdot \Phi(s_{k_0} + z) \rangle \\
&= \langle \eta_0, s_{k_0} + z \rangle - \frac{1}{p} W_p(\langle \eta_0, s_{k_0} + z \rangle) \\
&= \langle \eta_0, s_{k_0} + z \rangle - \frac{1}{p} W_p(\langle \eta_0, s_{k_0} + z \rangle)
\end{aligned}$$

Projecting to the  $+$ -component for  $W_p$  we obtain

$$\psi(0) = \left(1 - \frac{1}{p}\right) \cdot \langle \eta_0, s_{k_0} + z \rangle^+.$$

On the other hand, we have

$$\begin{aligned}
\langle \eta_0, s_{k_0} + z \rangle^+ &= \frac{1}{2} \cdot \left( \langle \eta_0, s_{k_0} + z \rangle + W_p(\langle \eta_0, s_{k_0} + z \rangle) \right) \\
&= \frac{1}{2} \cdot \left( \langle \eta_0, s_{k_0} + z \rangle - \langle W_p(\eta_0), W_p(s_{k_0} + z) \rangle \right) \\
&= \frac{1}{2} \cdot \left( \langle \eta_0, s_{k_0} + z \rangle + \langle \eta_0, W_p(s_{k_0} + z) \rangle \right) \\
&= \frac{1}{2} \cdot \langle \eta_0, (s_{k_0} + z) + W_p(s_{k_0} + z) \rangle \\
&= \frac{1}{2} \cdot \langle \eta_0, s_{k_0} + W_p(s_{k_0}) \rangle \\
&= \frac{1}{2} \cdot \langle \eta_0, \sigma(f) \rangle.
\end{aligned}$$

Finally, we use Proposition 3.1(2) to conclude that  $\langle \eta_0, \sigma(f) \rangle = -\frac{1}{2} \cdot \sigma(f)$ . Hence  $\psi(0) = \frac{1}{4} \left(1 - \frac{1}{p}\right) \sigma(f)$  and Lemma 5.3 is proved  $\square$

**Proof of Theorem B.** Combining Lemma 5.1 and Lemma 5.3 we obtain

$$-2 \cdot u'(k_0) \cdot \rho(f) = \sigma(f).$$

Hence  $\mathcal{L}(f) = -2 \cdot u'(k_0) = -2 \cdot p^{-\frac{k_0}{2}} \cdot \alpha'(k_0)$  and Theorem B is proved.  $\square$

## REFERENCES

- [1] Coleman, R. F.: Classical and overconvergent modular forms. *Invent. Math.* 124 (1996), no. 1-3, 215–241.
- [2] Coleman, R. F.:  $p$ -adic Banach spaces and families of modular forms. *Invent. Math.* 127 (1997), no. 3, 417–479.
- [3] Coleman, R. F.: A  $p$ -adic Shimura isomorphism and  $p$ -adic periods of modular forms.  $p$ -adic monodromy and the Birch and Swinnerton-Dyer conjecture (Boston, MA, 1991), 21–51, *Contemp. Math.*, 165, Amer. Math. Soc., Providence, RI, 1994.
- [4] Coleman, R. F.: Reciprocity laws on curves. *Compositio Math.* 72 (1989), no. 2, 205–235.
- [5] Coleman, R. F.; Iovita, A.: Hidden structures on semistable curves. preprint 2007.

- [6] Coleman, R. F.; Stevens, G.; Teitelbaum, J.: Numerical experiments on families of  $p$ -adic modular forms. Computational perspectives on number theory (Chicago, IL, 1995), 143–158, AMS/IP Stud. Adv. Math., 7, Amer. Math. Soc., Providence, RI, 1998.
- [7] Coleman, R. F.; Teitelbaum, J.: Numerical solution of the  $p$ -adic hypergeometric equation.  $p$ -adic monodromy and the Birch and Swinnerton-Dyer conjecture (Boston, MA, 1991), 53–62, Contemp. Math., 165, Amer. Math. Soc., Providence, RI, 1994.
- [8] Colmez, P.: Invariants  $\mathcal{L}$  et dérivées de valeurs propres de Frobenius.
- [9] Colmez, Pierre: La conjecture de Birch et Swinnerton-Dyer  $p$ -adique. Séminaire Bourbaki 2002-03, exp. 919, Astérisque No. 294 (2004), ix, 251–319.
- [10] Colmez, P.: Zéros supplémentaires de fonctions  $L$   $p$ -adiques de formes modulaires. Algebra and Number Theory, edited by R. Tandon, Hindustan book agency (2005), 193–210.
- [11] Greenberg, R.; Stevens, G.: On the conjecture of Mazur, Tate, and Teitelbaum.  $p$ -adic monodromy and the Birch and Swinnerton-Dyer conjecture (Boston, MA, 1991), 183–211, Contemp. Math., 165, Amer. Math. Soc., Providence, RI, 1994.
- [12] Greenberg, R.; Stevens, G.:  $p$ -adic  $L$ -functions and  $p$ -adic periods of modular forms. Invent. Math. 111 (1993), no. 2, 407–447.
- [13] Kato, K.; Kurihara, M.; Tsuji, T.: cours au centre Émile Borel, premier semestre 1997.
- [14] Katz, N.:  $p$ -adic properties of modular forms. Modular Functions of one variable III, SLN 350 (1972), 69–190.
- [15] Mazur, B.: On monodromy invariants occurring in global arithmetic, and Fontaine's theory.  $p$ -adic monodromy and the Birch and Swinnerton-Dyer conjecture (Boston, MA, 1991), 1–20, Contemp. Math., 165, Amer. Math. Soc., Providence, RI, 1994.
- [16] Mazur, B.; Tate, J.; Teitelbaum, J.: On  $p$ -adic analogues of the conjectures of Birch and Swinnerton-Dyer. Invent. Math. 84 (1986), no. 1, 1–48.
- [17] Stevens, G.: cours au centre Émile Borel, premier semestre 2000.
- [18] Teitelbaum, J.: Values of  $p$ -adic  $L$ -functions and a  $p$ -adic Poisson kernel. Invent. Math. 101 (1990), no. 2, 395–410.