
Professor Jennifer Balakrishnan, jbala@bu.edu

What is on today

1	Cramer's Rule, Volume, and Linear Transformations	1
2	Vector Spaces and Subspaces	2

1 Cramer's Rule, Volume, and Linear Transformations

Lay–Lay–McDonald §3.3 pp. 179 – 186

Today we give a geometric interpretation of the determinant.

Theorem 1. *If A is a 2×2 matrix, the area of the parallelogram determined by the columns of A is $|\det A|$. If A is a 3×3 matrix, the volume of the parallelepiped determined by the columns of A is $|\det A|$.*

Example 2. *Calculate the area of the parallelogram determined by the points $(-2, -2)$, $(0, 3)$, $(4, -1)$, and $(6, 4)$.*

Determinants can be used to describe an important geometric property of linear transformations in the plane and in \mathbb{R}^3 . If T is a linear transformation and S is a set in the domain of T , let $T(S)$ denote the set of images of points in S . We are interested in how the area (or volume) of $T(S)$ compares with the area (or volume) of the original set S .

Theorem 3. *Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation determined by a 2×2 matrix A . If S is a parallelogram in \mathbb{R}^2 , then*

$$\text{area of } T(S) = |\det A| \cdot \text{area of } S.$$

If T is determined by a 3×3 matrix A , and if S is a parallelepiped in \mathbb{R}^3 , then

$$\text{volume of } T(S) = |\det A| \cdot \text{volume of } S.$$

It turns out that the conclusions of the above theorem hold whenever S is a region in \mathbb{R}^2 with finite area or a region in \mathbb{R}^3 with finite volume.

Example 4. Let a and b be positive numbers. Find the area of the region E bounded by the ellipse whose equation is

$$\frac{x_1^2}{a^2} + \frac{x_2^2}{b^2} = 1.$$

2 Vector Spaces and Subspaces

Lay–Lay–McDonald §4.1 pp. 192 – 197

The work we've been doing with vectors in \mathbb{R}^n can be understood in a more general framework once we have the notion of a *vector space*, which will be our object of study today.

Definition 5. A (real) vector space is a nonempty set V of objects, called vectors, on which are defined two operations, addition and multiplication by scalars (real numbers), subject to the ten axioms below. The axioms must hold for all vectors $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$ and for all scalars c, d .

1. $\mathbf{u} + \mathbf{v} \in V$.
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$.
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$.
4. There is a zero vector $\mathbf{0} \in V$ such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$.
5. For each $\mathbf{u} \in V$, there is a vector $-\mathbf{u} \in V$ such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$.
6. $c\mathbf{u} \in V$.
7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$.
8. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$.
9. $c(d\mathbf{u}) = (cd)\mathbf{u}$.
10. $1\mathbf{u} = \mathbf{u}$.

Note that the zero vector $\mathbf{0}$ is unique, and for each $\mathbf{u} \in V$, its negative $-\mathbf{u}$ is unique.

Example 6. *The spaces \mathbb{R}^n for $n \geq 1$ are vector spaces.*

Example 7. *For $n \geq 0$, the set P_n of polynomials of degree at most n consists of all polynomials of the form*

$$p(t) = a_0 + a_1t + \cdots + a_nt^n,$$

where the coefficients a_0, a_1, \dots, a_n are real numbers. If $p(t) = a_0 \neq 0$, the degree of p is zero. If all of the coefficients are zero, p is called the zero polynomial. Show that P_n is a vector space.

Example 8. *Let V be the set of all real-valued functions defined on a set D (where D is \mathbb{R} or some interval on the real line). Show that V is a vector space.*

In many problems, a vector space consists of a subset of vectors from some larger vector space. In this case, only three of the ten vector space axioms need to be checked; the rest are automatically satisfied.

Definition 9. *A subspace of a vector space V is a subset H of V that has three properties:*

- 1. The zero vector of V is in H .*
- 2. H is closed under vector addition: $\mathbf{u}, \mathbf{v} \in H \Rightarrow \mathbf{u} + \mathbf{v} \in H$.*
- 3. H is closed under multiplication by scalars: if c is a scalar and $\mathbf{u} \in H$, then $c\mathbf{u} \in H$.*

Example 10. *Is the set consisting of the zero vector in a vector space V a subspace of V ?*

Example 11. Let P be the set of all polynomials with real coefficients, with the usual operations in P . Then P is a subspace of the space of all real-valued functions on \mathbb{R} . Also, for each $n \geq 0$, P_n is a subspace of P .

Example 12. The vector space \mathbb{R}^2 is not a subspace of \mathbb{R}^3 , since \mathbb{R}^2 is not a subset of \mathbb{R}^3 . However, the set $H = \left\{ \begin{bmatrix} s \\ t \\ 0 \end{bmatrix} \right\}$ where $s, t \in \mathbb{R}$ is a subset of \mathbb{R}^3 . Show that H is a subspace of \mathbb{R}^3 .

Example 13. Consider a plane in \mathbb{R}^3 not through the origin. Is it a subspace of \mathbb{R}^3 ?

Example 14. Let V be a vector space, and let $\mathbf{v}_1, \mathbf{v}_2 \in V$. Let $H = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}$ be the set of all linear combinations of $\mathbf{v}_1, \mathbf{v}_2$. Show that H is a subspace of V .

The argument in the previous example can be generalized to prove the following:

Theorem 15. If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are in a vector space V , then $\text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a subspace of V .

We call $\text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ the subspace spanned (or generated) by $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$. Given any subspace H of V , a spanning (or generating) set for H is a set $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ in H such that $H = \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$.

Example 16. Let H be the set of all vectors of the form $(a - 3b, b - a, a, b)$, where a, b are arbitrary real numbers. Show that H is a subspace of \mathbb{R}^4 .

Example 17. For what values of h will \mathbf{y} be in the subspace of \mathbb{R}^3 spanned by $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ if

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 5 \\ -4 \\ -7 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} -3 \\ 1 \\ 0 \end{bmatrix}, \mathbf{y} = \begin{bmatrix} -4 \\ 3 \\ h \end{bmatrix} ?$$

Example 18. Show that the set H of all points of \mathbb{R}^2 of the form $(3a, 2 + 5a)$ is not a vector space.