

## Multivariable chain rules

Today we discuss two types of chain rules. Both generalize the single variable chain rule.

## Chain Rule—Type I

Consider a vector-valued function  $\mathbf{P}(t) = x(t)\mathbf{i} + y(t)\mathbf{j}$  which parameterizes a curve in the  $xy$ -plane and a function  $f(x, y)$ . How is the derivative  $df/dt$  related to the partial derivatives of  $f(x, y)$  and the derivative  $\mathbf{P}'(t)$ ?

**Example.** Consider  $\mathbf{P}(t) = t\mathbf{i} + t^2\mathbf{j}$  and  $f(x, y) = 2x^2 + y^2$ .

**Chain Rule.** The derivative of the composition  $f(\mathbf{P}(t))$  is given by

$$\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}.$$

This version of the Chain Rule has an important formulation in terms of the gradient of  $f$ .

**Definition.** Given a function  $f(x, y)$  that is differentiable at the point  $(a, b)$ . Then the *gradient vector* of  $f$  at  $(a, b)$  is the vector

$$\nabla f(a, b) = \frac{\partial f}{\partial x}(a, b)\mathbf{i} + \frac{\partial f}{\partial y}(a, b)\mathbf{j}.$$

Sometimes the gradient vector of  $f$  is denoted  $\mathbf{grad}f(a, b)$ .

**Restatement of the Chain Rule.** The derivative of the composition  $f(\mathbf{P}(t))$  is

$$\left. \frac{df}{dt} \right|_{t=t_0} = \nabla f(\mathbf{P}(t_0)) \cdot \mathbf{P}'(t_0).$$

**Example.** We return to  $\mathbf{P}(t) = t\mathbf{i} + t^2\mathbf{j}$  and  $f(x, y) = 2x^2 + y^2$ .

Animation of this chain rule

**Example.** Use the polar curve  $r = \cos 2\theta$  to parameterize a curve  $\mathbf{P}(t)$  in the  $xy$ -plane and consider the composition  $f(\mathbf{P}(t))$  where

$$f(x, y) = y^2 - x^2.$$

This chain rule has some important theoretical implications as well.

**Theorem.**

1. Let  $f(x, y)$  be a differentiable function such that  $\nabla f(x, y) = \mathbf{0}$  for all  $(x, y)$ . Then  $f(x, y)$  is a constant function.
2. If  $g(x, y)$  and  $h(x, y)$  are two differentiable functions such that

$$\nabla g(x, y) = \nabla h(x, y)$$

for all  $(x, y)$ . Then  $g(x, y) = h(x, y) + K$  for some constant  $K$ .

## Chain Rule—Type II

For this situation, consider a function  $f(x, y)$  of two variables and suppose that the variables  $x$  and  $y$  are functions of other variables.

For example, consider  $x$  and  $y$  as a function of the polar coordinates  $r$  and  $\theta$ . That is,

$$x = r \cos \theta \quad \text{and} \quad y = r \sin \theta.$$

**Example.** Let  $f(x, y) = xy + y^2$ . What is the angular rate of change of  $f(x, y)$  at the point  $(x, y) = (1, 2)$ ?