

Multivariable chain rules

Today we discuss two types of chain rules. Both generalize the single variable chain rule.

Chain Rule—Type I

Consider a vector-valued function $\mathbf{P}(t) = x(t)\mathbf{i} + y(t)\mathbf{j}$ which parameterizes a curve in the xy -plane and a function $f(x, y)$. How is the derivative df/dt related to the partial derivatives of $f(x, y)$ and the derivative $\mathbf{P}'(t)$?

Example. Consider $\mathbf{P}(t) = t\mathbf{i} + t^2\mathbf{j}$ and $f(x, y) = 2x^2 + y^2$.

Chain Rule. The derivative of the composition $f(\mathbf{P}(t))$ is given by

$$\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}.$$

This version of the Chain Rule has an important formulation in terms of the gradient of f .

Definition. Given a function $f(x, y)$ that is differentiable at the point (a, b) . Then the *gradient vector* of f at (a, b) is the vector

$$\nabla f(a, b) = \frac{\partial f}{\partial x}(a, b) \mathbf{i} + \frac{\partial f}{\partial y}(a, b) \mathbf{j}.$$

Sometimes the gradient vector of f is denoted $\mathbf{grad}f(a, b)$.

Restatement of the Chain Rule. The derivative of the composition $f(\mathbf{P}(t))$ is

$$\left. \frac{df}{dt} \right|_{t=t_0} = \nabla f(\mathbf{P}(t_0)) \cdot \mathbf{P}'(t_0).$$

Example. We return to $\mathbf{P}(t) = t\mathbf{i} + t^2\mathbf{j}$ and $f(x, y) = 2x^2 + y^2$.

Animation of this chain rule

Example. Use the polar curve $r = \cos 2\theta$ to parameterize a curve $\mathbf{P}(t)$ in the xy -plane and consider the composition $f(\mathbf{P}(t))$ where

$$f(x, y) = y^2 - x^2.$$

This chain rule has some important theoretical implications as well.

Theorem.

1. Let $f(x, y)$ be a differentiable function such that $\nabla f(x, y) = \mathbf{0}$ for all (x, y) . Then $f(x, y)$ is a constant function.
2. If $g(x, y)$ and $h(x, y)$ are two differentiable functions such that

$$\nabla g(x, y) = \nabla h(x, y)$$

for all (x, y) . Then $g(x, y) = h(x, y) + K$ for some constant K .

Chain Rule—Type II

For this situation, consider a function $f(x, y)$ of two variables and suppose that the variables x and y are functions of other variables.

For example, consider x and y as a function of the polar coordinates r and θ . That is,

$$x = r \cos \theta \quad \text{and} \quad y = r \sin \theta.$$

Example. Let $f(x, y) = xy + y^2$. What is the angular rate of change of $f(x, y)$ at the point $(x, y) = (1, 2)$?