


More about lines

Last class we used vector addition and scalar multiplication to describe a line in space. Given two points $A = (a_1, a_2, a_3)$ and $B = (b_1, b_2, b_3)$ on the line, we first calculate a direction vector \mathbf{D} for the line. It is

$$\mathbf{D} = (b_1 - a_1)\mathbf{i} + (b_2 - a_2)\mathbf{j} + (b_3 - a_3)\mathbf{k}.$$

If \mathbf{P}_1 is a position vector representing any point on the line, then a (position) vector equation for the line is

$$\mathbf{L}(t) = \mathbf{P}_1 + t\mathbf{D}.$$



Example. Find a vector equation for the line ℓ that contains the points $(1, 1, 2)$ and $(3, 7, -2)$.

Summary: A direction vector for ℓ is $\mathbf{D} = 2\mathbf{i} + 6\mathbf{j} - 4\mathbf{k}$.

Vector form for ℓ :

$$\mathbf{L}(t) = (1 + 2t)\mathbf{i} + (1 + 6t)\mathbf{j} + (2 - 4t)\mathbf{k}$$

Parametric form for ℓ :

$$\begin{cases} x(t) = 1 + 2t \\ y(t) = 1 + 6t \\ z(t) = 2 - 4t \end{cases}$$

Symmetric form for ℓ :

$$\frac{x - 1}{2} = \frac{y - 1}{6} = \frac{z - 2}{-4}$$

Here's another example that has a slightly different symmetric form:

Example. Find an equation of the line through $(2, 4, 6)$ and $(1, 6, 6)$.

Dot Product

Given two vectors \mathbf{A} and \mathbf{B} , their **dot product** is defined to be

$$\mathbf{A} \cdot \mathbf{B} = |\mathbf{A}| |\mathbf{B}| \cos \theta,$$

where θ is the angle between \mathbf{A} and \mathbf{B} .

Important: Note that we start with two vectors and end up with a scalar. Therefore, we cannot take the dot product of three vectors in a row. Before we do a few examples, I would like to make a few observations.

1. Suppose $|\mathbf{A}|, |\mathbf{B}| \neq 0$. Then the two vectors \mathbf{A} and \mathbf{B} are perpendicular (orthogonal) $\iff \mathbf{A} \cdot \mathbf{B} = 0$.
2. There is a close relationship between projections and the dot product.

$$\text{comp}_{\mathbf{A}}(\mathbf{B}) = |\mathbf{B}| \cos \theta = \frac{\mathbf{A} \cdot \mathbf{B}}{|\mathbf{A}|} \quad \text{proj}_{\mathbf{A}}(\mathbf{B}) = (|\mathbf{B}| \cos \theta) \frac{\mathbf{A}}{|\mathbf{A}|} = \frac{(\mathbf{A} \cdot \mathbf{B})}{|\mathbf{A}|^2} \mathbf{A}$$

3. We can derive an algebraic formula for the dot product using the law of cosines. Let $\mathbf{C} = \mathbf{B} - \mathbf{A}$.

The law of cosines is

$$|\mathbf{C}|^2 = |\mathbf{A}|^2 + |\mathbf{B}|^2 - 2|\mathbf{A}||\mathbf{B}| \cos \theta.$$

Therefore,

$$\begin{aligned} 2(\mathbf{A} \cdot \mathbf{B}) &= |\mathbf{A}|^2 + |\mathbf{B}|^2 - |\mathbf{C}|^2 \\ &= a_1^2 + a_2^2 + a_3^2 + b_1^2 + b_2^2 + b_3^2 - (b_1 - a_1)^2 - (b_2 - a_2)^2 - (b_3 - a_3)^2. \end{aligned}$$

So $2(\mathbf{A} \cdot \mathbf{B}) = 2(a_1b_1 + a_2b_2 + a_3b_3)$, and therefore,

$$\mathbf{A} \cdot \mathbf{B} = a_1b_1 + a_2b_2 + a_3b_3.$$

This equation gives us an algebraic formula with geometric applications.

Examples. Let $\mathbf{A} = 2\mathbf{i} + \mathbf{j} + \mathbf{k}$ and $\mathbf{B} = 3\mathbf{i} + 4\mathbf{j} + 3\mathbf{k}$.

1. Compute the angle between \mathbf{A} and \mathbf{B} .

2. Compute the length of the projection of \mathbf{A} in the direction of \mathbf{B} .

Now let's use the dot product to derive the triangle inequality. Recall that

$$|\mathbf{A}|^2 = a_1^2 + a_2^2 + a_3^2,$$

so

$$|\mathbf{A}|^2 = \mathbf{A} \cdot \mathbf{A}.$$

We can therefore derive

$$\begin{aligned} |\mathbf{A} + \mathbf{B}|^2 &= (\mathbf{A} + \mathbf{B}) \cdot (\mathbf{A} + \mathbf{B}) \\ &= \mathbf{A} \cdot \mathbf{A} + 2(\mathbf{A} \cdot \mathbf{B}) + \mathbf{B} \cdot \mathbf{B} \\ &= |\mathbf{A}|^2 + 2|\mathbf{A}||\mathbf{B}|\cos\theta + |\mathbf{B}|^2 \\ &\leq |\mathbf{A}|^2 + 2|\mathbf{A}||\mathbf{B}| + |\mathbf{B}|^2 \\ &\leq (|\mathbf{A}| + |\mathbf{B}|)^2 \end{aligned}$$

and arrive at triangle inequality

$$|\mathbf{A} + \mathbf{B}| \leq |\mathbf{A}| + |\mathbf{B}|.$$

The length function $|\mathbf{A}|$ has four important properties:

1. $|\mathbf{A}| \geq 0$ for all \mathbf{A}
2. $|\mathbf{A}| = 0 \iff \mathbf{A} = \mathbf{0}$
3. $|r\mathbf{A}| = |r||\mathbf{A}|$
4. $|\mathbf{A} + \mathbf{B}| \leq |\mathbf{A}| + |\mathbf{B}|$

The dot product has the following properties:

1. $\mathbf{A} \cdot \mathbf{B} = \mathbf{B} \cdot \mathbf{A}$
2. $\mathbf{A} \cdot \mathbf{A} = |\mathbf{A}|^2$
3. $r(\mathbf{A} \cdot \mathbf{B}) = (r\mathbf{A}) \cdot \mathbf{B} = \mathbf{A} \cdot r\mathbf{B}$
4. $\mathbf{A} \cdot (\mathbf{B} + \mathbf{C}) = \mathbf{A} \cdot \mathbf{B} + \mathbf{A} \cdot \mathbf{C}$
5. $|\mathbf{A} \cdot \mathbf{B}| \leq |\mathbf{A}||\mathbf{B}|$ (which follows from $\mathbf{A} \cdot \mathbf{B} = |\mathbf{A}||\mathbf{B}| \cos \theta$)

Now let's apply what we have learned to a problem involving lines.

Example. Let ℓ_1 be line through $(1, 0, 0)$ and $(1, 2, 2)$. Let ℓ_2 be the line through $(1, 1, 1)$ and $(1 + \sqrt{2}, 2, 2)$. Do these lines intersect? What is the angle of intersection?