

Limits and continuity

In order to be able to do calculus for multivariable functions, we need to be able to talk about limits.

Informal definition. We say that

$$\lim_{(x,y) \rightarrow (a,b)} f(x,y) = L$$


if $f(x,y) \rightarrow L$ as $(x,y) \rightarrow (a,b)$ along all paths in the xy -plane.

Here are two examples to illustrate some of the issues that arise.

Example. Consider the limit of the function

$$f(x,y) = \frac{\sin(x^2 + y^2)}{x^2 + y^2}$$


as $(x,y) \rightarrow (0,0)$. The graph of $f(x,y)$ for $(x,y) \neq (0,0)$ is



Example. Consider the limit of the function

$$g(x, y) = \frac{2xy}{x^2 + y^2}$$

as $(x, y) \rightarrow (0, 0)$. The graph of $g(x, y)$ for $(x, y) \neq (0, 0)$ is




Partial derivatives

Consider a function of two variables $f(x, y)$. How do we talk about its rate of change at a given point?

Definition. The partial derivative of $f(x, y)$ in the x -direction at the point (x_0, y_0) is defined by

$$\frac{\partial f}{\partial x}(x_0, y_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h, y_0) - f(x_0, y_0)}{h}.$$

We vary x but keep y constant as we take the limit.



Example. Consider $f(x, y) = 9 - x^2 - y^2$. Let's calculate

$$\frac{\partial f}{\partial x}(1, 2)$$

directly from this definition.

There is another, more efficient way to calculate this partial derivative.

Let's try a more complicated example.

Example. Consider $g(x, y) = y \ln(xy) + y$.

The partial derivative with respect to y is defined in a similar fashion.

Definition. The partial derivative of $f(x, y)$ in the y -direction at the point (x_0, y_0) is defined by

$$\frac{\partial f}{\partial y}(x_0, y_0) = \lim_{h \rightarrow 0} \frac{f(x_0, y_0 + h) - f(x_0, y_0)}{h}.$$

We keep x constant and vary y as we take the limit.

Example. Consider $g(x, y) = y \ln(xy) + y$ again and calculate $\partial g / \partial y$ this time.