

More on the example from last class

**Example.** Once again consider


$$\frac{d\mathbf{Y}}{dt} = \begin{pmatrix} -3 & 1 \\ -1 & 0 \end{pmatrix} \mathbf{Y}.$$

For this example, the eigenvalues are

$$\lambda = \frac{-3 \pm \sqrt{5}}{2}.$$

Both are negative.

The slope of the eigenline that corresponds to the “fast” eigenvalue  $\lambda_1 = \frac{1}{2}(-3 - \sqrt{5})$  is approximately 0.4, and the slope of the eigenline that corresponds to the “slow” eigenvalue  $\lambda_2 = \frac{1}{2}(-3 + \sqrt{5})$  is approximately 2.6.




Sketching component graphs

Once we understand the phase portrait, we should also be able to sketch the component graphs without `HPGSystemSolver`.

Let's sketch the  $x(t)$ - and  $y(t)$ -graphs that correspond to the initial conditions  $(-3, 2)$  and  $(3, 2)$  for

$$\frac{d\mathbf{Y}}{dt} = \begin{pmatrix} -3 & 1 \\ -1 & 0 \end{pmatrix} \mathbf{Y}.$$




Case 2:  $\lambda_1 < 0 < \lambda_2$ .

We did this case at the start of last class.

**Example.** Consider


$$\frac{d\mathbf{Y}}{dt} = \begin{pmatrix} 4 & -5 \\ -2 & 1 \end{pmatrix} \mathbf{Y}.$$




Case 3:  $0 < \lambda_1 < \lambda_2$ .

**Example.** Consider


$$\frac{d\mathbf{Y}}{dt} = \begin{pmatrix} 3 & -1 \\ 1 & 0 \end{pmatrix} \mathbf{Y}.$$




Summary for real and distinct (nonzero) eigenvalues



sink ( $\lambda_1 < \lambda_2 < 0$ )



saddle ( $\lambda_1 < 0 < \lambda_2$ )



source ( $0 < \lambda_1 < \lambda_2$ )

## Complex eigenvalues

What happens if the eigenvalues of the system are complex numbers?

**Example.** Consider

$$\frac{d\mathbf{Y}}{dt} = \begin{pmatrix} -3 & 2 \\ -1 & -1 \end{pmatrix} \mathbf{Y}.$$

Let's see that happens if we take a look at this system using `MatrixFields` and then we'll compute the eigenstuff for this matrix.

Eigenvalues:

Eigenvectors:

(Lots of blank space on the next page.)

We now have a complex-valued solution of the form

$$\mathbf{Y}_c(t) = e^{(-2+i)t} \begin{pmatrix} 2 \\ 1+i \end{pmatrix}.$$

We are interested in real-valued solutions. What good is this complex-valued solution?