

More on Laplace transforms

Definition. The *Laplace transform* of the function $y(t)$ is the function

$$Y(s) = \int_0^{\infty} y(t) e^{-st} dt.$$

This transform is an “operator” (a function on functions). It transforms the function $y(t)$ into the function $Y(s)$.

For example, last class we showed that

$$\int_0^{\infty} e^{-st} dt = \begin{cases} \frac{1}{s}, & \text{if } s > 0; \\ \text{undefined,} & \text{if } s \leq 0. \end{cases}$$

This is the Laplace transform of the function that is constantly 1.

Notation: We often represent this operator using the script letter \mathcal{L} . In other words,

$$\mathcal{L}[y] = Y.$$

For example,

$$\mathcal{L}[1] = \frac{1}{s}.$$

Note that, even if $y(t)$ is defined for all t , the Laplace transform $Y(s)$ may not be defined for all s .

Example. Let's compute $\mathcal{L}[e^{at}]$ using the definition and the improper integrals we have already computed:

Examples. Using *Mathematica* to calculate the improper integrals, we see that:

$$\mathcal{L}[\sin t] = \frac{1}{s^2 + 1} \quad \text{for } s > 0,$$

$$\mathcal{L}[e^{2t} \sin 3t] = \frac{3}{s^2 - 4s + 13} \quad \text{for } s > 2$$

$$\mathcal{L}[t^4] = \frac{24}{s^5} \quad \text{for } s > 0$$

$$\mathcal{L}[\sin 2t] = \frac{2}{s^2 + 4} \quad \text{for } s > 0,$$

$$\mathcal{L}[t \cos \sqrt{2}t] = \frac{s^2 - 2}{(s^2 + 2)^2} \quad \text{for } s > 0$$

$$\mathcal{L}[e^{i\omega t}] = \frac{1}{s - i\omega} \quad \text{for } s > 0$$

Properties of the Laplace transform There are two properties of the Laplace transform that make it well suited for solving linear differential equations:

1. $\mathcal{L} \left[\frac{dy}{dt} \right] = s\mathcal{L}[y] - y(0)$
2. \mathcal{L} is a linear transform

Both of these properties are extremely important, but the surprising one is #1. Let's consider

$$\mathcal{L} \left[\frac{dy}{dt} \right] = \int_0^{\infty} \left(\frac{dy}{dt} \right) e^{-st} dt.$$

In fact, before we consider the improper integral, let's apply the method of integration by parts to the indefinite integral

$$\int \left(\frac{dy}{dt} \right) e^{-st} dt.$$

(Blank space on the top of the next page.)

Now let's see how we can use the Laplace transform to solve an initial-value problem.

Example. Solve the IVP

$$\frac{dy}{dt} - 3y = e^{2t}, \quad y(0) = 4.$$

