

More on the kernel and range of a linear transformation

Recall that differentiation is a linear transformation if we do a good job of identifying the vector spaces involved.

Example. Let V_1 be the vector space of all continuously differentiable functions $f : \mathbb{R} \rightarrow \mathbb{R}$ and let V_2 be the vector space of all continuous functions $f : \mathbb{R} \rightarrow \mathbb{R}$. The operation of differentiation is a linear transformation from V_1 to V_2 . That is, the transformation $D : V_1 \rightarrow V_2$ given by

$$D(f) = f'$$

is a linear transformation.

Example. What are the kernel and the range of the differentiation transformation D mentioned above?

Bases for vector spaces and subspaces

Given a vector space or subspace V , we often find it convenient to express it as the span of a few vectors. A basis for V is a spanning set that contains as few vectors as possible.

Definition. A set of vectors $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ is a *basis* for V if

1. it is linearly independent, and
2. it spans V .

Example. The standard basis $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ of \mathbb{R}^n .

Example. The two vectors

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

form a basis of \mathbb{R}^2 .

Example. The set $\{x^3, x^2, x, 1\}$ is a basis of \mathbb{P}_3 .

Example. The set $\{x^3, x^3 + x^2, x, 1\}$ is another basis of \mathbb{P}_3 .

We need ways of determining bases of vector spaces and their subspaces. The “casting-out procedure” produces a basis from a spanning set.

The casting-out procedure

Given a vector subspace S spanned by $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$, we can obtain a basis B for S by casting out the vectors that are linear combinations of the preceding vectors. More precisely, let

1. $B_1 = \{\mathbf{v}_1\}$ as long as $\mathbf{v}_1 \neq \mathbf{0}$, and
2. for $i \geq 2$,
 - (a) (cast out) $B_i = B_{i-1}$ if \mathbf{v}_i is in $\text{Span } B_{i-1}$, or
 - (b) (keep) $B_i = B_{i-1} \cup \{\mathbf{v}_i\}$ if \mathbf{v}_i is not in $\text{Span } B_{i-1}$.

Then the final result B_k is a basis B for S .

Example. Let’s apply the casting-out procedure to the set

$$\{x^3 + 1, x, x^2, x^2 - x, 4, x^3\}$$

of polynomials in \mathbb{P}_3 .

Theorem. (similar to The Spanning Set Theorem, Lay, p. 239) Let $S = \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$. Then the final result B_k of the casting-out procedure applied to $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ is a basis for S .

The proof of the casting-out procedure is posted on the web site, and we will not discuss it in class. However, to understand the proof of the theorem, it is helpful to consider the example above along with the sets B_1, B_2, \dots, B_6 . We get

$$B_1 = \{x^3 + 1\}$$

$$B_2 = \{x^3 + 1, x\}$$

$$B_3 = \{x^3 + 1, x, x^2\}$$

$$B_4 = B_3$$

$$B_5 = \{x^3 + 1, x, x^2, 4\}$$

$$B_6 = B_5.$$

Bases for Nul \mathbf{A} and Col \mathbf{A}

Last class we did an example that showed how we can produce a basis for Nul \mathbf{A} .

Example. Find a basis for the column space of

$$\mathbf{A} = \begin{bmatrix} 1 & -2 & 0 & 1 \\ 0 & 0 & 3 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Example. Find a basis for the column space of

$$\mathbf{B} = \begin{bmatrix} 1 & -2 & 0 & 1 \\ -1 & 2 & 3 & 1 \\ 0 & 0 & -3 & -2 \end{bmatrix}.$$

Fact: Suppose that \mathbf{A} and \mathbf{B} are row equivalent matrices. Then the linear dependence relations among the columns of \mathbf{A} are the same as the linear dependence relations among the columns of \mathbf{B} .

Why?

Warning: If you row reduce a matrix \mathbf{A} to a matrix \mathbf{B} in row echelon form, you identify the pivot columns of \mathbf{A} . To find a basis for $\text{Col } \mathbf{A}$, use the pivot columns of \mathbf{A} . **Do not use the pivot columns of \mathbf{B} .** Row reduction usually changes the column space of a matrix.