

More on subspaces of vector spaces

**Definition.** A nonempty subset  $S$  of a vector space  $V$  is a *subspace* of  $V$  if

1. the zero vector  $\mathbf{0}$  is in  $S$ ,
2. (closure under vector addition) for each  $\mathbf{v}_1$  and  $\mathbf{v}_2$  in  $S$ , the vector sum  $\mathbf{v}_1 + \mathbf{v}_2$  is in  $S$ , and
3. (closure under scalar multiplication) for each  $r$  in  $\mathbb{R}$  and each  $\mathbf{v}$  in  $S$ , the scalar multiple  $r\mathbf{v}$  is in  $S$ .

**Note.** A subspace  $S$  of a vector space  $V$  is a vector space in its own right.

**Example.** Consider the line  $x_2 = 3x_1$  in the vector space  $\mathbb{R}^2$ .

**Example.** Consider the line  $x_2 = x_1 + 1$  in the vector space  $\mathbb{R}^2$ .

**Example.** Let  $\mathbb{P}$  represent the vector space of all polynomial functions as discussed last class. Is  $\mathbb{P}$  a subspace of the vector space of all functions  $f : \mathbb{R} \rightarrow \mathbb{R}$ ?

**Example.** Consider the subset  $S = \text{Span}\{x, x^2\}$  within  $\mathbb{P}$ . Is  $S$  a subspace of  $\mathbb{P}$ ?

**Theorem.** If  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$  are vectors in a vector space  $V$ , then  $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$  is a subspace of  $V$ .



Subspaces associated to a matrix

There are three important subspaces associated to an  $m \times n$  matrix  $\mathbf{A}$ . Let  $\mathbf{c}_1, \dots, \mathbf{c}_n$  represent the columns of  $\mathbf{A}$ . That is,

$$\mathbf{A} = \left[ \begin{array}{c|c|c|c} \mathbf{c}_1 & \mathbf{c}_2 & \dots & \mathbf{c}_n \end{array} \right].$$

These column vectors are vectors in  $\mathbb{R}^m$ .

Let  $\mathbf{r}_1, \dots, \mathbf{r}_m$  represent the rows of  $\mathbf{A}$ . That is,

$$\mathbf{A} = \left[ \begin{array}{c} \mathbf{r}_1 \\ \mathbf{r}_2 \\ \vdots \\ \mathbf{r}_m \end{array} \right].$$

These row vectors are vectors in  $\mathbb{R}^n$ .

**The column space of  $\mathbf{A}$ .** The column space of  $\mathbf{A}$  is the span of the columns of  $\mathbf{A}$ . We write

$$\text{Col } \mathbf{A} = \text{Span}\{\mathbf{c}_1, \dots, \mathbf{c}_n\}.$$

**The row space of  $\mathbf{A}$ .** The row space of  $\mathbf{A}$  is the span of the rows of  $\mathbf{A}$ . We write

$$\text{Row } \mathbf{A} = \text{Span}\{\mathbf{r}_1, \dots, \mathbf{r}_m\}.$$

**The null space of  $\mathbf{A}$ .** The null space of  $\mathbf{A}$  is the set of all vectors  $\mathbf{x}$  in  $\mathbb{R}^n$  such that

$$\mathbf{A}\mathbf{x} = \mathbf{0}.$$

The null space of  $\mathbf{A}$  is denoted by  $\text{Nul } \mathbf{A}$ .

---

---

**Theorem.** Let  $\mathbf{A}$  be an  $m \times n$  matrix. The column space of  $\mathbf{A}$  is a subspace of  $\mathbb{R}^m$ , and the null space and the row space of  $\mathbf{A}$  are subspaces of  $\mathbb{R}^n$ .

**Application.** Any plane through the origin in  $\mathbb{R}^3$  is a subspace of  $\mathbb{R}^3$ .