

Subspaces associated to a matrix

There are three important subspaces associated to an  $m \times n$  matrix  $\mathbf{A}$ . Let  $\mathbf{c}_1, \dots, \mathbf{c}_n$  represent the columns of  $\mathbf{A}$ . That is,

$$\mathbf{A} = \left[ \begin{array}{c|c|c|c} \mathbf{c}_1 & \mathbf{c}_2 & \dots & \mathbf{c}_n \end{array} \right].$$

These column vectors are vectors in  $\mathbb{R}^m$ .

Let  $\mathbf{r}_1, \dots, \mathbf{r}_m$  represent the rows of  $\mathbf{A}$ . That is,

$$\mathbf{A} = \left[ \begin{array}{c} \mathbf{r}_1 \\ \mathbf{r}_2 \\ \vdots \\ \mathbf{r}_m \end{array} \right].$$

These row vectors are vectors in  $\mathbb{R}^n$ .

**The column space of  $\mathbf{A}$ .** The column space of  $\mathbf{A}$  is the span of the columns of  $\mathbf{A}$ . We write

$$\text{Col } \mathbf{A} = \text{Span}\{\mathbf{c}_1, \dots, \mathbf{c}_n\}.$$

**The row space of  $\mathbf{A}$ .** The row space of  $\mathbf{A}$  is the span of the rows of  $\mathbf{A}$ . We write

$$\text{Row } \mathbf{A} = \text{Span}\{\mathbf{r}_1, \dots, \mathbf{r}_m\}.$$

**The null space of  $\mathbf{A}$ .** The null space of  $\mathbf{A}$  is the set of all vectors  $\mathbf{x}$  in  $\mathbb{R}^n$  such that

$$\mathbf{A}\mathbf{x} = \mathbf{0}.$$

The null space of  $\mathbf{A}$  is denoted by  $\text{Nul } \mathbf{A}$ .

---

---

**Theorem.** Let  $\mathbf{A}$  be an  $m \times n$  matrix. The column space of  $\mathbf{A}$  is a subspace of  $\mathbb{R}^m$ , and the null space and the row space of  $\mathbf{A}$  are subspaces of  $\mathbb{R}^n$ .

**Application.** Any plane through the origin in  $\mathbb{R}^3$  is a subspace of  $\mathbb{R}^3$ .

**Example.** Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & -2 & 0 & 4 & 0 \\ 0 & 0 & 1 & -9 & 0 \\ 2 & -4 & 0 & 8 & 1 \end{bmatrix}.$$

Express the null space of  $\mathbf{A}$  as the span of as few vectors as possible.

The consistency of a system of linear equations can be viewed as a statement about the column space of the coefficient matrix.

**Fact.** The linear system  $\mathbf{Ax} = \mathbf{b}$  is consistent if and only if  $\mathbf{b}$  is an element of the column space of  $\mathbf{A}$ .

Here is how Lay (p. 232) contrasts  $\text{Nul } \mathbf{A}$  and  $\text{Col } \mathbf{A}$  for an  $m \times n$  matrix  $\mathbf{A}$ :

Nul $\mathbf{A}$	Col $\mathbf{A}$
1. Nul $\mathbf{A}$ is a subspace of $\mathbb{R}^n$ .	1. Col $\mathbf{A}$ is a subspace of $\mathbb{R}^m$ .
2. Nul $\mathbf{A}$ is implicitly defined; that is, you are given only a condition ( $\mathbf{Ax} = \mathbf{0}$ ) that vectors in Nul $\mathbf{A}$ must satisfy.	2. Col $\mathbf{A}$ is explicitly defined; that is, you are told how to build vectors in Col $\mathbf{A}$ .
3. It takes time to find vectors in Nul $\mathbf{A}$ . Row operations on $[\mathbf{A} \ \mathbf{0}]$ are required.	3. It is easy to find vectors in Col $\mathbf{A}$ . The columns of $\mathbf{A}$ are displayed; others are formed from them.
4. There is no obvious relation between Nul $\mathbf{A}$ and the entries in $\mathbf{A}$ .	4. There is an obvious relation between Col $\mathbf{A}$ and the entries in $\mathbf{A}$ , since each column of $\mathbf{A}$ is in Col $\mathbf{A}$ .
5. A typical vector $\mathbf{v}$ in Nul $\mathbf{A}$ has the property that $\mathbf{Av} = \mathbf{0}$ .	5. A typical vector $\mathbf{v}$ in Col $\mathbf{A}$ has the property that the equation $\mathbf{Ax} = \mathbf{v}$ is consistent.
6. Given a specific vector $\mathbf{v}$ , it is easy to tell if $\mathbf{v}$ is in Nul $\mathbf{A}$ . Just compute $\mathbf{Av}$ .	6. Given a specific vector $\mathbf{v}$ , it may take time to tell if $\mathbf{v}$ is in Col $\mathbf{A}$ . Row operations on $[\mathbf{A} \ \mathbf{v}]$ are required.
7. Nul $\mathbf{A} = \{\mathbf{0}\}$ if and only if the equation $\mathbf{Ax} = \mathbf{0}$ has only the trivial solution.	7. Col $\mathbf{A} = \mathbb{R}^m$ if and only if the equation $\mathbf{Ax} = \mathbf{b}$ has a solution for every $\mathbf{b}$ in $\mathbb{R}^m$ .
8. Nul $\mathbf{A} = \{\mathbf{0}\}$ if and only if the linear transformation $\mathbf{x} \mapsto \mathbf{Ax}$ is one-to-one.	8. Col $\mathbf{A} = \mathbb{R}^m$ if and only if the linear transformation $\mathbf{x} \mapsto \mathbf{Ax}$ maps $\mathbb{R}^n$ onto $\mathbb{R}^m$ .

Item 8 in both lists suggest two subspaces that are intimately connected with any linear transformation from one vector space to another.

**Definition.** A transformation  $L : V_1 \rightarrow V_2$  from a vector space  $V_1$  to a vector space  $V_2$  is linear if

1.  $L(\mathbf{v}_1 + \mathbf{v}_2) = L(\mathbf{v}_1) + L(\mathbf{v}_2)$  for all vectors  $\mathbf{v}_1$  and  $\mathbf{v}_2$  in  $V_1$ , and
2.  $L(r\mathbf{v}) = rL(\mathbf{v})$  for all  $\mathbf{v}$  in  $V_1$  and all  $r$  in  $\mathbb{R}$ .

**Example.** Let  $V_1$  be the vector space of all continuously differentiable functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  and let  $V_2$  be the vector space of all continuous functions  $f : \mathbb{R} \rightarrow \mathbb{R}$ . The operation of differentiation is a linear transformation from  $V_1$  to  $V_2$ . That is, the transformation  $D : V_1 \rightarrow V_2$  given by

$$D(f) = f'$$

is a linear transformation.

Associated to any linear transformation are two important subspaces.

**Definition.** The kernel of  $L : V_1 \rightarrow V_2$  is the subset of  $V_1$  given by

$$\{\mathbf{v}_1 \mid L(\mathbf{v}_1) = \mathbf{0}\}.$$

The range of  $L$  is the subset of  $V_2$  given by

$$\{\mathbf{v}_2 \mid L(\mathbf{v}_1) = \mathbf{v}_2 \text{ for some } \mathbf{v}_1 \text{ in } V_1\}.$$

**Fact.** Both the kernel and the range of a linear transformation are subspaces. The kernel is a subspace of  $V_1$ , and the range is a subspace of  $V_2$ .

For a matrix transformation from  $\mathbb{R}^n$  to  $\mathbb{R}^m$  determined by the matrix  $\mathbf{A}$ , its range is  $\text{Col } \mathbf{A}$ , and its kernel is  $\text{Nul } \mathbf{A}$ .

**Example.** What are the kernel and range of the transformation  $p : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  determined by the matrix

$$\frac{1}{3} \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix} ?$$

(This example was first introduced on September 21.)